

CS 441 Discrete Mathematics for CS
Lecture 21b

Relations

Milos Hauskrecht
milos@cs.pitt.edu
5329 Sennott Square

Cartesian product (review)

Let $A = \{a_1, a_2, \dots, a_k\}$ and $B = \{b_1, b_2, \dots, b_m\}$.

The Cartesian product $A \times B$ is defined by a set of pairs
 $\{(a_1, b_1), (a_1, b_2), \dots, (a_1, b_m), \dots, (a_k, b_m)\}$.

Cartesian product defines a product set, or a set of all ordered arrangements of elements in sets in the Cartesian product.

Binary relation

Definition: Let A and B be two sets. A **binary relation from A to B** is a subset of a Cartesian product $A \times B$.

- Let $R \subseteq A \times B$ means R is a set of ordered pairs of the form (a,b) where $a \in A$ and $b \in B$.
- We use the notation **$a R b$ to denote $(a,b) \in R$** and **$a \not R b$ to denote $(a,b) \notin R$** . If **$a R b$** , we say a is related to b by R.

Example: Let $A = \{a,b,c\}$ and $B = \{1,2,3\}$.

- Is $R = \{(a,1), (b,2), (c,2)\}$ a relation from A to B? **Yes.**
- Is $Q = \{(1,a), (2,b)\}$ a relation from A to B? **No.**
- Is $P = \{(a,a), (b,c), (b,a)\}$ a relation from A to A? **Yes**


Representing binary relations

- We can graphically represent a binary relation R as follows:
 - if **$a R b$** then draw an arrow from a to b.

$$a \rightarrow b$$

Example:

- Let $A = \{0, 1, 2\}$, $B = \{u,v\}$ and $R = \{(0,u), (0,v), (1,v), (2,u)\}$
- Note: $R \subseteq A \times B$.
- **Graph:**



Representing binary relations

- We can represent a binary relation R by a **table** showing (marking) the ordered pairs of R .

Example:

- Let $A = \{0, 1, 2\}$, $B = \{u, v\}$ and $R = \{(0, u), (0, v), (1, v), (2, u)\}$


- **Table:**

R	u	v	or	R	u	v
0	x	x		0	1	1
1		x		1	0	1
2	x			2	1	0

Relations and functions

- Relations represent **one to many relationships** between elements in A and B .


- **Example:**




- What is the difference between a **relation** and a **function** from A to B ?

Relations and functions

- Relations represent **one to many relationships** between elements in A and B.
- **Example:**



- What is the difference between a **relation and a function from A to B**? A function defined on sets A,B $A \rightarrow B$ assigns to each element in the domain set A exactly one element from B. So it is a **special relation**.



Relation on the set

Definition: A relation on the set A is a relation from A to itself.

Example 1:

- Let $A = \{1,2,3,4\}$ and $R_{\text{div}} = \{(a,b) \mid a \text{ divides } b\}$
- What does R_{div} consist of?
- $R_{\text{div}} = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,4), (3,3), (4,4)\}$

R	1	2	3	4
1	x	x	x	x
2		x		x
3			x	
4				x

Relation on the set

Example:

- Let $A = \{1,2,3,4\}$.
- Define $a R_{\neq} b$ if and only if $a \neq b$.

$$R_{\neq} = \{(1,2), (1,3), (1,4), (2,1), (2,3), (2,4), (3,1), (3,2), (3,4), (4,1), (4,2), (4,3)\}$$

- | R | 1 | 2 | 3 | 4 |
|---|---|---|---|---|
| 1 | | x | x | x |
| 2 | x | | x | x |
| 3 | x | x | | x |
| 4 | x | x | x | |

Binary relations

- **Theorem:** The number of binary relations on a set A , where $|A| = n$ is:

$$2^{n^2}$$

- **Proof:**
- If $|A| = n$ then the cardinality of the Cartesian product $|A \times A| = n^2$.
- R is a binary relation on A if $R \subseteq A \times A$ (that is, R is a subset of $A \times A$).
- The number of subsets of a set with k elements : 2^k
- The number of subsets of $A \times A$ is : $2^{|A \times A|} = 2^{n^2}$

Binary relations

- **Example:** Let $A = \{1,2\}$
 - What is $A \times A = \{(1,1),(1,2),(2,1),(2,2)\}$
 - **List of possible relations (subsets of $A \times A$):**

• \emptyset	1	}	16
• $\{(1,1)\} \quad \{(1,2)\} \quad \{(2,1)\} \quad \{(2,2)\}$	4		
• $\{(1,1), (1,2)\} \quad \{(1,1),(2,1)\} \quad \{(1,1),(2,2)\}$	6		
• $\{(1,2),(2,1)\} \quad \{(1,2),(2,2)\} \quad \{(2,1),(2,2)\}$				
• $\{(1,1),(1,2),(2,1)\} \quad \{(1,1),(1,2),(2,2)\}$	4		
• $\{(1,1),(2,1),(2,2)\} \quad \{(1,2),(2,1),(2,2)\}$				
• $\{(1,1),(1,2),(2,1),(2,2)\}$	1		
- Use formula: $2^4 = 16$

Properties of relations

Definition (reflexive relation) : A relation R on a set A is called **reflexive** if $(a,a) \in R$ for every element $a \in A$.

Example 1:

- Assume relation $R_{\text{div}} = \{(a,b), \text{ if } a \mid b\}$ on $A = \{1,2,3,4\}$
- **Is R_{div} reflexive?**
- $R_{\text{div}} = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,4), (3,3), (4,4)\}$
- **Answer:** Yes. $(1,1), (2,2), (3,3),$ and $(4,4) \in A$.

Reflexive relation

Reflexive relation

- $R_{\text{div}} = \{(a \mid b), \text{ if } a \mid b\}$ on $A = \{1,2,3,4\}$
- $R_{\text{div}} = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,4), (3,3), (4,4)\}$

$$\text{MR}_{\text{div}} = \begin{array}{cccc} & 1 & 2 & 3 & 4 \\ \begin{array}{c} 1 \\ 2 \\ 3 \\ 4 \end{array} & 0 & 1 & 0 & 1 \\ & 0 & 0 & 1 & 0 \\ & 0 & 0 & 0 & 1 \end{array}$$

- **A relation R is reflexive** if and only if MR has 1 in every position on its main diagonal.

Properties of relations

Definition (reflexive relation) : A relation R on a set A is called **reflexive** if $(a,a) \in R$ for every element $a \in A$.

Example 2:

- Relation R_{fun} on $A = \{1,2,3,4\}$ defined as:
 - $R_{\text{fun}} = \{(1,2), (2,2), (3,3)\}$.
- **Is R_{fun} reflexive?**
- **No.** It is not reflexive since $(1,1) \notin R_{\text{fun}}$.

Properties of relations

Definition (irreflexive relation): A relation R on a set A is called **irreflexive** if $(a,a) \notin R$ for every $a \in A$.

Example 1:

- Assume relation R_{\neq} on $A = \{1,2,3,4\}$, such that $\mathbf{a R_{\neq} b}$ if and only if $a \neq b$.
- **Is R_{\neq} irreflexive?**
- $R_{\neq} = \dots$

Properties of relations

Definition (irreflexive relation): A relation R on a set A is called **irreflexive** if $(a,a) \notin R$ for every $a \in A$.

Example 1:

- Assume relation R_{\neq} on $A = \{1,2,3,4\}$, such that $\mathbf{a R_{\neq} b}$ if and only if $a \neq b$.
- **Is R_{\neq} irreflexive?**
- $R_{\neq} = \{(1,2), (1,3), (1,4), (2,1), (2,3), (2,4), (3,1), (3,2), (3,4), (4,1), (4,2), (4,3)\}$
- **Answer:** Yes. Because $(1,1), (2,2), (3,3)$ and $(4,4) \notin R_{\neq}$

Irreflexive relation

Irreflexive relation

- R_{\neq} on $A = \{1, 2, 3, 4\}$, such that $\mathbf{a R_{\neq} b}$ if and only if $a \neq b$.
- $R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}$

$$\text{MR} = \begin{array}{cccc} & 0 & 1 & 1 & 1 \\ & 1 & 0 & 1 & 1 \\ & 1 & 1 & 0 & 1 \\ & 1 & 1 & 1 & 0 \end{array}$$

- **A relation R is irreflexive** if and only if MR has 0 in every position on its main diagonal.

Properties of relations

Definition (irreflexive relation): A relation R on a set A is called **irreflexive** if $(a, a) \notin R$ for every $a \in A$.

Example 2:

- R_{fun} on $A = \{1, 2, 3, 4\}$ defined as:
 - $R_{\text{fun}} = \{(1, 2), (2, 2), (3, 3)\}$.
- **Is R_{fun} irreflexive?**
- **Answer: No.** Because $(2, 2)$ and $(3, 3) \in R_{\text{fun}}$

Properties of relations

Definition (symmetric relation): A relation R on a set A is called **symmetric** if

$$\forall a, b \in A \quad (a,b) \in R \rightarrow (b,a) \in R.$$

Example 1:

- $R_{\text{div}} = \{(a,b) \mid a \mid b\}$ on $A = \{1,2,3,4\}$
- **Is R_{div} symmetric?**
- $R_{\text{div}} = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,4), (3,3), (4,4)\}$
- **Answer: No.** It is not symmetric since $(1,2) \in R$ but $(2,1) \notin R$.

Properties of relations

Definition (symmetric relation): A relation R on a set A is called **symmetric** if

$$\forall a, b \in A \quad (a,b) \in R \rightarrow (b,a) \in R.$$

Example 2:

- R_{\neq} on $A = \{1,2,3,4\}$, such that **$a R_{\neq} b$** if and only if $a \neq b$.
- **Is R_{\neq} symmetric?**
- $R_{\neq} = \{(1,2), (1,3), (1,4), (2,1), (2,3), (2,4), (3,1), (3,2), (3,4), (4,1), (4,2), (4,3)\}$
- **Answer: Yes.** If $(a,b) \in R_{\neq} \rightarrow (b,a) \in R_{\neq}$

Symmetric relation

Symmetric relation:

- R_{\neq} on $A = \{1,2,3,4\}$, such that $\mathbf{a} R_{\neq} \mathbf{b}$ if and only if $a \neq b$.
- $R_{\neq} = \{(1,2), (1,3), (1,4), (2,1), (2,3), (2,4), (3,1), (3,2), (3,4), (4,1), (4,2), (4,3)\}$

$$\text{MR} = \begin{array}{cccc} & 0 & 1 & 1 & 1 \\ & 1 & 0 & 1 & 1 \\ & 1 & 1 & 0 & 1 \\ & 1 & 1 & 1 & 0 \end{array}$$

- A relation R is symmetric if and only if $m_{ij} = m_{ji}$ for all i, j .

Properties of relations

Definition (symmetric relation): A relation R on a set A is called symmetric if

$$\forall a, b \in A \quad (a,b) \in R \rightarrow (b,a) \in R.$$

Example 3:

- Relation R_{fun} on $A = \{1,2,3,4\}$ defined as:
 - $R_{\text{fun}} = \{(1,2), (2,2), (3,3)\}$.
- Is R_{fun} symmetric?
- **Answer: No.** For $(1,2) \in R_{\text{fun}}$ there is no $(2,1) \in R_{\text{fun}}$